

Happy Numbers II - Trial

The process of “**breaking**” an integer is defined as summing the squares of its digits. For example, the result of breaking the integer **125** is $(1^2 + 2^2 + 5^2) = 30$. An integer **N** is **happy** if after “**breaking**” it repeatedly the result reaches 1. If the result never reaches 1 no matter how many times the “**breaking**” is repeated, then **N** is not a happy number.

TASK

Write a program that given an integer **T** (number of test cases) and **T** integers, determines for each number whether it is a happy number or not.

CONSTRAINTS

$$1 \leq T \leq 1,080,000$$

$$2 \leq N \leq 2,147,483,647 \text{ (number for determining whether it is happy or not)}$$

Input

- The first line contains an integer **T**.
- next 1...T lines contain an integer **N** for determining whether it is happy or not.

Output

- T lines containing a single integer **N** which is the number of times the process had to be done to determine that **N** is happy, or **-1** if **N** is not happy.

Example

Input:

2
19
204

Output:

4
-1

- 1) 19 : $1^2 + 9^2 = 82$
- 2) 82 : $8^2 + 2^2 = 68$
- 3) 68 : $6^2 + 8^2 = 100$
- 4) 100 : $1^2 + 0^2 + 0^2 = 1$

The solution for 19 is 4 because we discovered that the integer 19 is happy after we repeated the process 4 times.

204 → 20 → 4 → 16 → 37 → 58 → 89 → 145 → 42 → 20 → 4 → 16 → 37 → 58 → 89 → 145
.....

204 is not a happy number because after breaking it several times the results start repeating so we can deduce that if we continue breaking it, the result will never reach 1.