

Internally Stable Sets

A weighted finite undirected graph is a triple $G = (V, E, w)$ consisting of vertex set V , edge set $E \subseteq \{(u, v) : u, v \in V, u \neq v\}$, and vertex weighting function w such that $w(u) \geq 0, \forall u \in V$ and $w(K) = \sum_{u \in K} w(u), K \subseteq V$. For $u \in V$ and $K \subseteq V$, $N(u)$ and $N(K)$ will denote the neighboring vertex sets of u and K respectively, formally defined as:

$$N(u) = \{v : \{u, v\} \in E\}, N(K) = \bigcup_{u \in K} N(u)$$

A vertex set $K \subseteq V$ satisfying $N(K) \cap K = \emptyset$ is called *internally stable* (also known as independent or anti-clique). In this problem you must find an internally stable set B such that $w(B) = \max\{w(S)\}$, where S belongs to the set of all internally stable sets of that graph.

Input

t – the number of test cases [$t \leq 100$]

n k – [n – number of vertices ($2 \leq n \leq 200$), k – number of edges ($1 \leq k \leq n*(n-1)/2$)]

then n numbers follows (w_i - the weight of i -th vertex) [$0 \leq w_i \leq 2^{31}-1$]

then k pairs of numbers follows denoting the edge between the vertices (s_i s_j edge between i -th and j -th vertices) [$1 \leq s_i, s_j \leq n$]

Output

For each test case output *MaxWeight* – the weight of a maximum internally stable set of the given graph. [$0 \leq \text{MaxWeight} \leq 2^{31}-1$]

Example

Input:

```
2
5 6
10 20 30 40 50
1 2
1 5
2 3
3 4
3 5
4 5

4 4
10 4 10 14
1 2
2 3
3 4
4 1
```

Output:

```
70
20
```